

AN INTERPRETATION OF THE LASCoux-LECLERC-THIBON ALGORITHM AND GRADED REPRESENTATION THEORY

ALEXANDER KLESHCHEV AND DAVID NASH

ABSTRACT. We use graded Specht modules to calculate the graded decomposition numbers for the Iwahori-Hecke algebra of the symmetric group over a field of characteristic zero at a root of unity. The algorithm arrived at is the Lascoux-Leclerc-Thibon algorithm in disguise. Thus we interpret the algorithm in terms of graded representation theory.

1. INTRODUCTION

Let H_d be the Hecke algebra of the symmetric group S_d defined over $\mathbb{C}$ with a parameter ξ which is a primitive e^{th} root of unity. The problem of finding dimensions of the irreducible H_d -modules is equivalent to the problem of finding decomposition numbers for H_d . This problem has been solved by Ariki [A] who proved a conjecture of Lascoux, Leclerc and Thibon [LLT] which gives an algorithm for computing the decomposition numbers.

Recently, Brundan and the first author [BK1, BK3] have defined *graded decomposition numbers* for H_d and proved a graded analogue of Ariki's theorem. The grading provides new information about H_d -modules, which is collected into a *graded character* of a module. Graded characters of Specht modules were computed in [BKW]. This together with the main results of [BK3] is sufficient to yield an algorithm for computing graded decomposition numbers for H_d .

Despite the different approach, the algorithm is equivalent to the one suggested in [LLT], although this equivalence is not immediately obvious. In section 5 we explain the equivalence of the two algorithms. Note that our approach gives a new interpretation of some coefficients computed in the LLT algorithm. We direct the reader to [L] for examples of such computations.

Acknowledgement. In the previous version of the paper we missed the equivalence of our algorithm and the LLT algorithm. We are grateful to the referee for pointing this out.

2. PRELIMINARIES

2.1. Basic objects. Let $\mathbb{F}$ be an algebraically closed field and $\xi \in \mathbb{F}^\times$. The *quantum characteristic*, e , is the smallest positive integer such that $1 + \xi + \xi^2 + \cdots + \xi^{e-1} = 0$, where we set $e := 0$ if no such integer exists. Set

Supported in part by the NSF grant DMS-0654147. The paper was completed while the authors were visiting the Isaac Newton Institute for Mathematical Sciences in Cambridge, U.K., whom we thank for hospitality and support.

$I := \mathbb{Z}/e\mathbb{Z}$. For any $i \in I$ we have a well-defined element

$$\nu(i) := \begin{cases} i & \text{if } \xi = 1, \\ \xi^i & \text{if } \xi \neq 1, \end{cases}$$

of $\mathbb{F}$. Throughout the paper, q is an indeterminate and we define the *bar-involution* on $\mathbb{Z}[q, q^{-1}]$ by $\overline{p(q)} = p(q^{-1})$ for all $p(q) \in \mathbb{Z}[q, q^{-1}]$. We set

$$[n]_q := \frac{q^n - q^{-n}}{q - q^{-1}} \text{ and } [n]_q^! := [n]_q [n-1]_q \cdots [1]_q.$$

The *Iwahori-Hecke algebra* of S_d with parameter ξ , is the $\mathbb{F}$ -algebra $H_d = H_d(\mathbb{F}, \xi)$ with generators $T_1, T_2, \dots, T_{d-1}$ and relations

$$\begin{aligned} T_r^2 &= (\xi - 1)T_r + \xi \quad (1 \leq r < d), \\ T_r T_{r+1} T_r &= T_{r+1} T_r T_{r+1} \quad (1 \leq r < d-1), \\ T_r T_s &= T_s T_r \quad (1 \leq r, s < d, |r-s| > 1). \end{aligned}$$

The group algebra $\mathbb{F}S_d$ of the symmetric group appears in the special case when $\xi = 1$. In this case $e = \text{char } \mathbb{F}$.

The *Jucys-Murphy elements* of H_d are:

$$L_r = \begin{cases} (1, r) + (2, r) + \cdots + (r-1, r) & \text{if } \xi = 1 \\ \xi^{1-r} T_{r-1} \cdots T_2 T_1 T_2 \cdots T_{r-1} & \text{if } \xi \neq 1 \end{cases} \quad (1 \leq r \leq d).$$

It follows from [G, Lemma 4.7] and [K, Lemma 7.1.2] that the eigenvalues of $L_1, \dots, L_d$ on any finite dimensional H_d -module are of the form $\nu(i)$ for $i \in I$. For $\mathbf{i} = (i_1, \dots, i_d) \in I^d$ and a finite dimensional H_d -module V , we define the $\mathbf{i}$ -*weight space* of V to be

$$V_{\mathbf{i}} := \{v \in V \mid (L_r - \nu(i_r))^N v = 0 \text{ for } N \gg 0 \text{ and } r = 1, \dots, d\}.$$

Then we have the *weight space decomposition* $V = \bigoplus_{\mathbf{i} \in I^d} V_{\mathbf{i}}$.

2.2. Partitions and tableaux. We use the partition notation from [BK3]. In particular, $\mathcal{P}_d$ is the set of all partitions of d . Given a partition $\lambda = (\lambda_1, \lambda_2, \dots) \in \mathcal{P}_d$, we define $\sigma_k(\lambda) = \sum_{i=1}^k \lambda_i$. Denote the usual *dominance order* on $\mathcal{P}_d$ by ' $\triangleright$ ', see [J1]. A partition $\lambda = (\lambda_1, \lambda_2, \dots)$ is called *e-restricted* if $\lambda_r - \lambda_{r+1} < e$ for all $r = 1, 2, \dots$. We let $\mathcal{RP}_d \subset \mathcal{P}_d$ be the subset of all *e-restricted* partitions of d .

Let $\lambda \in \mathcal{P}_d$. A node $A \in \lambda$ is called *removable* (for λ) if $\lambda \setminus \{A\}$ has the shape of a partition. A node $B \notin \lambda$ is called *addable* (for λ) if $\lambda \cup \{B\}$ has the shape of a partition. Given any set of removable nodes $\mathbf{A} = \{A_1, \dots, A_m\}$ for λ , we denote $\lambda_{\mathbf{A}} := \lambda \setminus \{A_1, A_2, \dots, A_m\}$.

If the node A is in row a and column b , we write $A = (a, b)$, and then the *residue* of A is defined to be $\text{res } A := b - a \pmod{e} \in I$. For $i \in I$, a node A is called an *i-node* if $\text{res } A = i$. For $\lambda, \mu \in \mathcal{P}_d$, we write $\lambda \sim \mu$ if and only if for each $i \in I$ the number of *i-nodes* in λ is equal to that in μ .

Given $\mu \in \mathcal{P}_d$ we call $\lambda \in \mathcal{P}_d$ a *move* for μ if $\lambda \trianglelefteq \mu$ and $\lambda \sim \mu$. We denote the set of moves for μ by $M(\mu)$. Put $M(\mu, \lambda) := \{\nu \in M(\mu) \mid \lambda \in M(\nu)\}$. Note that if λ is not a move of μ then $M(\mu, \lambda)$ is empty. For $\lambda \in M(\mu)$, we define the *distance* between λ and μ to be

$$l(\mu, \lambda) := \sum_{k \geq 1} \sigma_k(\mu) - \sigma_k(\lambda).$$

Since $\lambda \trianglelefteq \mu$, we know that $\sigma_k(\lambda) \leq \sigma_k(\mu)$ for all $k \geq 1$. So $l(\mu, \lambda) \geq 0$, with equality if and only if $\lambda = \mu$. Moreover, if $\nu \in M(\mu, \lambda)$, then $l(\mu, \nu) \leq l(\mu, \lambda)$ with equality if and only if $\nu = \lambda$.

Following [J2], for $m \in \mathbb{Z}_{>0}$, we define the m^{th} ladder L_m as the set of nodes of the form $(1 + k, m - k(e - 1))$ for all non-negative integers k with $k < m/(e - 1)$. Informally, the ladders are straight lines with slope $1/e$. Note that our ladders are transposed to those of James, since we are using the newer Dipper-James-Mathas notation for Specht modules. The nodes in the ladder L_m all have the residue $m - 1 \pmod{e}$, which we refer to as the *residue* of the ladder, and denote $\text{res } L_m$. For a partition λ and a positive integer m , we set $r_m(\lambda) := |\lambda \cap L_m|$. Denote by t_λ the maximal index such that $r_{t_\lambda}(\lambda) \neq 0$, and refer to the ladders $L_1, L_2, \dots, L_{t_\lambda}$ as the *ladders of λ* (some of them could have trivial intersection with λ). A ladder L_m is *bottom complete* for λ if whenever a node $A = (a, b) \in L_m$ belongs to λ , all other nodes $(a', b') \in L_m$ with $a' > a$ also belong to λ .

Lemma 2.1. [J2, 1.2] *Let $\lambda \in \mathcal{P}_d$. Then λ is e -restricted if and only if all ladders are bottom complete for λ .*

Given nodes $A = (a_1, a_2)$ and $B = (b_1, b_2)$ we say A is *above* (resp. *below*) B if $a_1 < b_1$ (resp. $a_1 > b_1$). Let $\lambda \in \mathcal{P}_d$. For a removable i -node A in λ we define the *degree* of A to be:

$$d_A(\lambda) := \#\{\text{addable } i\text{-nodes below } A\} \\ - \#\{\text{removable } i\text{-nodes below } A\}.$$

For $\lambda \in \mathcal{P}_d$, let $\mathcal{T}(\lambda)$ denote the set of all standard λ -tableaux. Let $T \in \mathcal{T}(\lambda)$. For $s \in \mathbb{Z}_{\geq 0}$ we denote by $T_{\leq s}$ and $\text{sh}(T_{\leq s})$ the tableau obtained by retaining the nodes of T labeled by the numbers $1, \dots, s$ and its shape respectively, so that $T_{\leq s} \in \mathcal{T}(\text{sh}(T_{\leq s}))$. Let A be the node of λ labeled by d in T . Following [BKW], we define the degree of T inductively by:

$$\deg(T) = \begin{cases} d_A(\lambda) + \deg(T_{\leq d-1}) & \text{if } d > 0, \\ 0 & \text{if } d = 0. \end{cases}$$

For $T \in \mathcal{T}(\lambda)$ we also have the associated *residue sequence*:

$$\mathbf{i}^T = (i_1, \dots, i_d) \in I^d,$$

where i_r is the residue of the node labeled by r in T for $1 \leq r \leq d$.

2.3. Representation theory of H_d . From the work of Dipper, James and Mathas [DJM], the algebra H_d has a special family of modules $\{S(\mu) \mid \mu \in \mathcal{P}_d\}$, labeled by the partitions of d , called *Specht modules*. This goes back to Dipper-James [DJ], although the Specht modules defined in [DJ] are different from those in [DJM]. Here we follow the conventions of [DJM].

If $e = 0$, then $\{S(\mu) \mid \mu \in \mathcal{P}_d\}$ is a complete irredundant set of the irreducible H_d -modules. In the more interesting case $e > 0$, the head $D(\mu)$ of $S(\mu)$ is irreducible, provided $\mu \in \mathcal{RP}_d$, and $\{D(\mu) \mid \mu \in \mathcal{RP}_d\}$ is a complete irredundant set of the irreducible H_d -modules.

In this paper, grading always means $\mathbb{Z}$ -grading. The algebra H_d has an explicit grading exhibited in [BK1]. Therefore we may speak of *graded H_d -modules*. In [BK3, §2.7], a notion of graded duality $\otimes$ on finite dimensional

graded H_d -modules is introduced. It is then shown that each irreducible H_d -module $D(\mu)$ has a unique grading which makes it into a *graded* H_d -module with $D(\mu)^\circledast \cong D(\mu)$. Moreover, by [BKW], the Specht modules are also gradable as H_d -modules in such a way that the natural projection $S(\mu) \twoheadrightarrow D(\mu)$ is a degree zero map whenever $\mu \in \mathcal{RP}_d$.

If $V = \bigoplus_{m \in \mathbb{Z}} V_m$ is a finite dimensional graded vector space, define $\text{qdim } V := \sum_{m \in \mathbb{Z}} (\dim V_m) q^m \in \mathbb{Z}[q, q^{-1}]$. Let $\mathcal{C}$ be the free $\mathbb{Z}[q, q^{-1}]$ -module on I^d . Given a finite dimensional graded H_d -module V , we define its *graded character* to be

$$\text{ch}_q V := \sum_{\mathbf{i} \in I^d} (\text{qdim } V_{\mathbf{i}}) \cdot \mathbf{i} \in \mathcal{C}$$

The graded characters of Specht modules are now as follows:

Theorem 2.2. [BKW, §4.3] *Let $\mu \in \mathcal{P}_d$. Then $\text{ch}_q S(\mu) = \sum_{\mathbf{T} \in \mathcal{T}(\mu)} q^{\deg(\mathbf{T})} \mathbf{i}^{\mathbf{T}}$.*

In particular, $\text{ch}_q S(\mu)$ depends only on μ and e . We extend the bar-involution from $\mathbb{Z}[q, q^{-1}]$ to $\mathcal{C}$ so that $\bar{\mathbf{i}} = \mathbf{i}$ for all $\mathbf{i} \in I^d$. The fact that $D(\mu)^\circledast \cong D(\mu)$ can now be interpreted as follows:

Theorem 2.3. [BK3, Theorem 4.18(3)] *Let $\lambda \in \mathcal{RP}_d$. Then $\text{ch}_q D(\lambda)$ is bar-invariant.*

When viewed as graded H_d -modules, the set $\{D(\mu) \mid \mu \in \mathcal{RP}_d\}$ forms a complete irredundant set of the irreducible graded H_d -modules *up to a grading shift*. Given $m \in \mathbb{Z}$ and a graded H_d -module V , we let $V\langle m \rangle$ denote the module obtained by shifting the grading of V up by m . Thus we have $\text{ch}_q V\langle m \rangle = q^m \text{ch}_q V$.

For $\mu \in \mathcal{P}_d$ and $\lambda \in \mathcal{RP}_d$, we define the corresponding *graded decomposition number* as follows

$$d_{\mu, \lambda} = d_{\mu, \lambda}(q) := \sum_{m \in \mathbb{Z}} a_m q^m \in \mathbb{Z}_{\geq 0}[q, q^{-1}],$$

where a_m is the multiplicity of $D(\lambda)\langle m \rangle$ in a graded composition series of $S(\mu)$. Note that $d_{\mu, \lambda}(1)$ is the usual decomposition number, so the following result easily follows from the well-known facts in the ungraded setting and the fact that the natural map $S(\mu) \twoheadrightarrow D(\mu)$ is of degree zero.

Theorem 2.4. *Let $\lambda \in \mathcal{RP}_d$ and $\mu \in \mathcal{P}_d$. Then*

- (i) $d_{\mu, \lambda} = 0$ unless $\lambda \in M(\mu)$.
- (ii) $d_{\lambda, \lambda} = 1$.

We have

$$\text{ch}_q S(\mu) = \sum_{\lambda \in \mathcal{RP}_d} d_{\mu, \lambda} \text{ch}_q D(\lambda).$$

By Theorem 2.4, the graded decomposition matrix $(d_{\mu, \lambda})$ is unitriangular, so the knowledge of the graded decomposition numbers implies the knowledge of the graded characters of the irreducible H_d -modules. The converse is also true since the graded characters of the irreducible H_d -modules are linearly independent, see e.g. [KL, Theorem 3.17]).

The following key fact is special for the case $\text{char } \mathbb{F} = 0$:

Theorem 2.5. [BK3, Theorem 3.9 and Corollary 5.15] *Let $\text{char } \mathbb{F} = 0$, $\lambda \in \mathcal{RP}_d$ and $\mu \in \mathcal{P}_d$. If $\mu \neq \lambda$, then $d_{\mu,\lambda} \in q\mathbb{Z}_{\geq 0}[q]$.*

3. THE LADDER WEIGHT

3.1. A dominance lemma. For $\lambda \in \mathcal{RP}_d$, let $t = t_\lambda$ be the index of its bottom ladder, and $r_t(\lambda) = |\lambda \cap L_t|$. Denote $\lambda \cap L_t = \{A_1, \dots, A_{r_t(\lambda)}\}$. Order the nodes of this set so that A_u is below A_s whenever $u < s$, and refer to the sequence $(A_1, \dots, A_{r_t(\lambda)})$ as the *bottom removable sequence* of λ . Observe that the nodes of the bottom removable sequence of λ are indeed removable nodes of λ . All nodes of this sequence have the same residue which we refer to as the *residue of the bottom removable sequence*.

The following technical result generalizes Lemma 1.1 in [KS].

Lemma 3.1. *Let $\lambda \in \mathcal{RP}_d$ and $\mu \in \mathcal{P}_d$, with $\lambda \not\leq \mu$. Let $\mathbf{A} = (A_1, \dots, A_r)$ be the bottom removable sequence for λ , and i be its residue. If $\mathbf{B} = \{B_1, \dots, B_r\}$ is any set of r removable i -nodes for μ then $\lambda \mathbf{A} \not\leq \mu \mathbf{B}$.*

Proof. Let $\lambda \mathbf{A} \leq \mu \mathbf{B}$. We need to show that $\lambda \leq \mu$. Let A_m be in row j_m of λ , and let B_m be in row l_m of μ for $m = 1, \dots, r$. By our convention $j_m = j_1 - (m - 1)$ for $1 \leq m \leq r$. We may also assume that $l_1 > \dots > l_r$.

Let $\lambda^m = \lambda \setminus \{A_{m+1}, \dots, A_r\}$ and $\mu^m = \mu \setminus \{B_{m+1}, \dots, B_r\}$ for $0 \leq m \leq r$. Then it suffices to show by induction on $m = 0, 1, \dots, r$ that $\lambda^m \leq \mu^m$, with the induction base case, $m = 0$, being our assumption.

Let $m > 0$ and assume by induction that $\lambda^{m-1} \leq \mu^{m-1}$. Note that

$$\begin{aligned} \sigma_k(\lambda^m) &= \begin{cases} \sigma_k(\lambda^{m-1}) & \text{if } k < j_m, \\ \sigma_k(\lambda^{m-1}) + 1 & \text{if } k \geq j_m, \end{cases} \\ \sigma_k(\mu^m) &= \begin{cases} \sigma_k(\mu^{m-1}) & \text{if } k < l_m, \\ \sigma_k(\mu^{m-1}) + 1 & \text{if } k \geq l_m. \end{cases} \end{aligned}$$

Since $\lambda^{m-1} \leq \mu^{m-1}$, we deduce that

$$\sigma_k(\mu^m) \geq \sigma_k(\lambda^m) \quad (\text{for } k \geq l_m \text{ or } k < j_m).$$

Observe that since $\mathbf{A}$ was the bottom removable sequence, it follows that j_1 is the bottom non-empty row in λ . Since λ is e -restricted, we know also that $\text{res}(j_1 + 1, 1) \neq i$. Furthermore, we have that row $j_1 + 1$ is empty in $\lambda_{\mathbf{A}}$, which implies that it is empty in $\mu_{\mathbf{B}}$ as well since $\lambda_{\mathbf{A}} \leq \mu_{\mathbf{B}}$.

Since B_1 is an addable i -node for $\mu_{\mathbf{B}}$, it follows that $l_1 < j_1 + 1$. Since $j_m = j_1 - (m - 1)$ and $l_n < l_{n-1}$ for all $1 \leq n \leq r$, it follows that $l_m \leq j_m$. Thus for all k we have either $k < j_m$ or $k \geq l_m$, which gives us that $\sigma_k(\mu^m) \geq \sigma_k(\lambda^m)$ for all $k > 0$. Hence $\mu^m \geq \lambda^m$ completing the proof. $\square$

3.2. Definition and properties of $\mathbf{j}^\lambda$. Let $\lambda \in \mathcal{RP}_d$ have bottom removable sequence $\mathbf{A} = (A_1, A_2, \dots, A_r)$ and define the *ladder weight* $\mathbf{j}^\lambda = (j_1, \dots, j_d)$ inductively as follows: $j_d = \text{res } A_r$ and $(j_1, \dots, j_{d-1}) = \mathbf{j}^{\lambda_{A_r}}$. The idea of the ladder weight appears in [LLT, §6.2].

Lemma 3.2. *Let $\lambda \in \mathcal{RP}_d$ and $\mu \in \mathcal{P}_d$, with $\mu \not\leq \lambda$. Then $\mathbf{j}^\lambda$ does not appear in $\text{ch}_q S(\mu)$. In particular, if μ is e -restricted, then $\mathbf{j}^\lambda$ does not appear in $\text{ch}_q D(\mu)$.*

Proof. We apply induction on d , the base case $d = 0$ being clear. Let $d > 0$ and suppose for a contradiction that $\mathbf{i}^T = \mathbf{j}^\lambda$ for some $T \in \mathcal{T}(\mu)$. Let $\mathbf{A} = (A_1, \dots, A_r)$ be the bottom removable sequence of λ and let i be its residue. Let $\mathbf{B} = \{B_1, \dots, B_r\}$ be the nodes of μ labeled in T with $d, d-1, \dots, d-r+1$. Since $\mathbf{i}^T = \mathbf{j}^\lambda$ we have that $\text{res } B_1 = \dots = \text{res } B_r = i$. Let $T' \in \mathcal{T}(\mu_{\mathbf{B}})$ be the tableau obtained from T by removing $B_1, \dots, B_r$. Then $\mathbf{i}^{T'} = \mathbf{j}^{\lambda_{\mathbf{A}}}$, whence $\mathbf{j}^{\lambda_{\mathbf{A}}}$ appears in $\text{ch}_q S(\mu_{\mathbf{B}})$. By the inductive assumption, $\mu_{\mathbf{B}} \supseteq \lambda_{\mathbf{A}}$. Now, by Lemma 3.1, $\mu \supseteq \lambda$, a contradiction. $\square$

Lemma 3.3. *Let $\lambda \in \mathcal{RP}_d$ and set $t = t_\lambda$, $r_m = r_m(\lambda)$, $R_m := r_1 + \dots + r_m$, and $\lambda(m) := \lambda \cap (L_1 \cup \dots \cup L_m)$ for $m > 0$. If $T \in \mathcal{T}(\lambda)$ has $\mathbf{i}^T = \mathbf{j}^\lambda$ then for each $m > 0$ we have $\text{sh}(T_{\leq R_m}) = \lambda(m)$.*

Proof. We apply induction on $m > 0$ with the induction base $m = 1$ being clear as $r_1 = 1$. Let $m > 1$ and assume that $\text{sh}(T_{\leq R_{m-1}}) = \lambda(m-1)$. Letting $\mathbf{B}$ denote the set of nodes in $\text{sh}(T_{\leq R_m}) \setminus \text{sh}(T_{\leq R_{m-1}})$, it suffices to prove that $\mathbf{B} = \lambda \cap L_m$. Since $|\mathbf{B}| = |\lambda \cap L_m| = r_m$, it is enough to prove that $\mathbf{B} \subseteq \lambda \cap L_m$. Observing that $\mathbf{B}$ is contained in λ it then remains to show that $\mathbf{B} \subseteq L_m$. Observe that the nodes of $\mathbf{B}$ must have residue $\text{res } L_m$ since $\mathbf{i}^T = \mathbf{j}^\lambda$. We know also that none of the nodes in $\mathbf{B}$ belong to any of the ladders $L_1, \dots, L_{m-1}$ since $\text{sh}(T_{\leq R_{m-1}}) = \lambda(m-1)$. We conclude that $\mathbf{B} \subseteq L_m$, completing the inductive step. $\square$

Let $\lambda \in \mathcal{P}_d$ and set $t = t_\lambda$ and $r_m = r_m(\lambda)$ for $m > 0$. We define

$$r_\lambda := [r_1]_q! [r_2]_q! \cdots [r_t]_q!.$$

Lemma 3.4. *If $\lambda \in \mathcal{RP}_d$ then $\mathbf{j}^\lambda$ has multiplicity r_λ in $\text{ch}_q S(\lambda)$.*

Proof. Assume that $\mathbf{i}^T = \mathbf{j}^\lambda$. Lemma 3.3 implies that for any tableau $T \in \mathcal{T}(\lambda)$ with $\mathbf{i}^T = \mathbf{j}^\lambda$, the numbers $d, d-1, \dots, d-r_t+1$ must appear in the bottom removable sequence $\mathbf{A} = \{A_1, \dots, A_{r_t}\}$ for λ . Moreover, all $r_t!$ possible permutations of those numbers occur. Since λ is e -restricted, by Lemma 2.1 all ladders are bottom complete for λ , and thus the possible labelings of $\mathbf{A}$ give a contribution of exactly $[r_t]_q!$ to the multiplicity of $\mathbf{j}^\lambda$ in $\text{ch}_q S(\lambda)$. The result follows by induction on d . $\square$

Corollary 3.5. *If $\lambda \in \mathcal{RP}_d$ then $\mathbf{j}^\lambda$ has multiplicity r_λ in $\text{ch}_q D(\lambda)$.*

Proof. By Lemma 3.4, $\mathbf{j}^\lambda$ appears in $\text{ch}_q S(\lambda)$ with multiplicity r_λ , and by Lemma 3.2, $\mathbf{j}^\lambda$ does not appear in $\text{ch}_q D(\mu)$ for $\mu \triangleleft \lambda$. Since composition factors of $S(\lambda)$ are of the form $D(\mu)$ for $\mu \leq \lambda$, the result follows. $\square$

In two special cases we can be more explicit.

Corollary 3.6. *Let λ be a partition of d with $\lambda_1 < e$. Then $\mathbf{j}^\lambda$ has multiplicity 1 in $\text{ch}_q S(\lambda)$ and in $\text{ch}_q D(\lambda)$.*

Proof. Observe that $\lambda_1 < e$ implies that $r_m \leq 1$ for all $1 \leq m \leq s_\lambda$. The result then follows directly from Lemma 3.4 and Corollary 3.5. $\square$

Corollary 3.7. *Let $\lambda = (l_1^{k_1}, l_2^{k_2}, \dots)$ be an e -restricted partition with $l_1 = e$ and $l_i > l_{i+1}$ for $i \geq 1$. Then the multiplicity of $\mathbf{j}^\lambda$ in $\text{ch}_q S(\lambda)$ and in $\text{ch}_q D(\lambda)$ is $([2]_q)^{k_1}$.*

Proof. Use Lemma 3.4 and Corollary 3.5 and the fact that λ has exactly k_1 ladders of size 2, the remaining ladders of λ having size at most 1. $\square$

3.3. Ladder weight multiplicity. Let V be a finite dimensional graded H_d -module. For any $\lambda \in \mathcal{RP}_d$ define $m_\lambda(V)$ to be the multiplicity of $\mathbf{j}^\lambda$ in $\text{ch}_q V$. We collect the important properties of the function m_λ :

Theorem 3.8. *Let $\lambda \in \mathcal{RP}_d$, $\mu \in \mathcal{P}_d$, and V be a finite dimensional graded H_d -module. Then:*

- (i) $m_\lambda(V) \in \mathbb{Z}_{\geq 0}[q, q^{-1}]$;
- (ii) if $m_\lambda(V) = 0$ then $[V : D(\lambda)] = 0$;
- (iii) $m_\lambda(S(\lambda)) = m_\lambda(D(\lambda)) = r_\lambda$;
- (iv) $m_\lambda(S(\mu)) = 0$ unless $\lambda \in M(\mu)$;
- (v) if $\mu \in \mathcal{RP}_d$, then $m_\lambda(D(\mu)) = 0$ unless $\lambda \in M(\mu)$;
- (vi) $m_\lambda(S(\mu)) = \sum_{\nu \in \mathcal{RP}_d \cap M(\mu, \lambda), \nu \neq \lambda} d_{\mu, \nu} m_\lambda(D(\nu)) + d_{\mu, \lambda} r_\lambda$.

Proof. (i) is clear from the definitions, and (iii) is a restatement of Lemma 3.4 and Corollary 3.5. Next, (ii) follows from (i) and (iii).

To see (iv), assume that $\lambda \notin M(\mu)$. Then either $\lambda \not\prec \mu$ or $\lambda \not\leq \mu$. In the first case it follows from Theorem 2.4 that $m_\lambda(S(\mu)) = 0$, and in the second case the same follows from Lemma 3.2. Now (v) follows from (iv) and (i).

Finally, (vi) follows from (i), (iii), (v), and Theorem 2.4 $\square$

4. THE ALGORITHM

4.1. Basic Algorithm. We give an easy algorithm to solve the following problem which will arise in the Main Algorithm below:

Problem 4.1. *Suppose $d(q) \in q\mathbb{Z}[q]$, and $m(q), r(q) \in \mathbb{Z}[q, q^{-1}]$ are such that $\overline{m(q)} = m(q)$, $\overline{r(q)} = r(q)$, and $r(q) \neq 0$. If $d(q)r(q) + m(q)$ is known and $r(q)$ is known, find $d(q)$ and $m(q)$.*

Remark 4.2. It is easy to see that Problem 4.1 has a unique solution.

We now explain an inductive algorithm to solve Problem 4.1. Clearly, if $d(q)r(q) + m(q) = 0$, then $d(q) = m(q) = 0$ by uniqueness. If $d(q)r(q) + m(q) \neq 0$ then we can write it in the form

$$d(q)r(q) + m(q) = \sum_{n=-N}^M a_n q^n \quad (\text{for } -N \leq M \text{ and } a_{-N} \neq 0, a_M \neq 0).$$

Note that the assumptions imply that $M \geq 0$ and $M \geq N$ (but we might have $N < 0$ if $m(q) = 0$). The algorithm proceeds by induction on the pairs of non-negative integers $(M, M+N)$ ordered lexicographically. The induction base is the pair $(0, 0)$ where $d(q) = 0$ and $m(q) = a_0$ by uniqueness. Let $(M, M+N) > (0, 0)$. This implies $M > 0$. We denote the top term of $r(q)$ by bq^R , and note that $R \geq 0$ since $r(q)$ is bar-invariant. We now consider two cases.

Case 1: $M > N$. As $m(q)$ is bar-invariant, the term $a_M q^M$ must come from $d(q)r(q)$. Thus $\frac{a_M}{b} q^{M-R}$ is a term in $d(q)$. Setting $d'(q) := d(q) -$

$\frac{a_M}{b}q^{M-R}$, we are reduced to solving the problem for $d'(q)r(q) + m(q)$ for which $M' < M$.

Case 2: $M = N$. As $d(q) \in q\mathbb{Z}[q]$, the term $a_{-N}q^{-N}$ must therefore come from $m(q)$. Since $m(q)$ is bar-invariant, $a_{-N}q^N$ must be a term in $m(q)$ also. Setting $m'(q) := m(q) - (a_{-N}q^{-N} + a_{-N}q^N)$, we are reduced to solving the problem for $d(q)r(q) + m'(q)$ for which $M' \leq M$ and $M' + N' < M + N$.

4.2. Main Algorithm. From now on we assume that $\text{char } \mathbb{F} = 0$. If $e = 0$, then the Specht modules are irreducible, so we also assume that we are in the interesting case $e > 0$, i.e we deal with the case of the Hecke algebra over a field of characteristic zero with parameter a primitive e^{th} root of unity. Under these assumptions, we will describe an algorithm for computing the graded decomposition numbers.

The algorithm relies heavily on Theorem 2.5, which is why we need the assumption $\text{char } \mathbb{F} = 0$.

Let $\mu \in \mathcal{P}_d$ and $\lambda \in \mathcal{RP}_d$. We will compute $d_{\mu,\lambda}$ by induction. However, this induction requires us to keep track of some extra information. By Theorem 2.4(i), $\lambda \notin M(\mu)$ implies $d_{\mu,\lambda} = 0$, so we assume $\lambda \in M(\mu)$. We now calculate $d_{\mu,\lambda}$ and $m_\lambda(D(\mu))$ by induction on the distance $l(\mu, \lambda)$ (of course, $m_\lambda(D(\mu))$ only makes sense when μ is e -restricted). Induction begins when $l(\mu, \lambda) = 0$, hence $\mu = \lambda$, and we have $d_{\mu,\mu} = 1$ by Theorem 2.4(ii) and $m_\mu(S(\mu)) = r_\mu$ by Theorem 3.8.

Let $l(\mu, \lambda) > 0$, so $\mu \triangleright \lambda$. By induction, we know the graded decomposition numbers $d_{\mu,\nu}$ for all $\nu \in \mathcal{RP}_d \cap M(\mu, \lambda)$ with $\nu \neq \lambda$ and the multiplicities $m_\lambda(D(\nu))$ for all $\nu \in \mathcal{RP}_d \cap M(\mu, \lambda)$ with $\nu \neq \mu$. To make the inductive step we need to compute $d_{\mu,\lambda}$ and, if μ is e -restricted, $m_\lambda(D(\mu))$.

If μ is not e -restricted, then by Theorem 3.8(vi), we have

$$d_{\mu,\lambda} = \frac{1}{r_\lambda} \left(m_\lambda(S(\mu)) - \sum_{\nu \in \mathcal{RP}_d \cap M(\mu, \lambda), \nu \neq \lambda} d_{\mu,\nu} m_\lambda(D(\nu)) \right),$$

where all the terms in the right hand side are known by induction and Theorem 2.2.

Let μ be e -restricted. By Theorem 3.8(vi), we have

$$m_\lambda(D(\mu)) + d_{\mu,\lambda} r_\lambda = m_\lambda(S(\mu)) - \sum_{\nu \in \mathcal{RP}_d \cap M(\mu, \lambda), \nu \neq \lambda, \nu \neq \mu} d_{\mu,\nu} m_\lambda(D(\nu)),$$

where all terms in the right hand side are known by induction and Theorem 2.2. Note r_λ is non-zero and bar-invariant, $d_{\mu,\lambda} \in q\mathbb{Z}_{\geq 0}[q]$ by Theorem 2.5, and $m_\lambda(D(\mu))$ is bar-invariant by Theorem 2.3. Hence, we are in the assumptions of Problem 4.1 with $m(q) = m_\lambda(D(\mu))$, $d(q) = d_{\mu,\lambda}$, and $r(q) = r_\lambda$. Now we apply the Basic Algorithm described in Section 4.1 to calculate $m_\lambda(D(\mu))$ and $d_{\mu,\lambda}$ and complete the inductive step.

Remark 4.3. In our algorithm, to calculate $d_{\mu,\lambda}$ and $m_\lambda(D(\mu))$ one only ever needs to compute $d_{\nu,\kappa}$ and $m_\kappa(D(\nu))$ for pairs (ν, κ) such that $l(\nu, \kappa) < l(\mu, \lambda)$ and $\nu, \kappa \in M(\mu, \lambda)$.

5. CONNECTION TO LLT

5.1. Grothendieck groups. Here we recast some notions from [LLT] and [BK3]. Let $[\text{Rep}(H_d)]$ denote the Grothendieck group of the category of finite dimensional graded H_d -modules. This is a free $\mathbb{Z}[q, q^{-1}]$ -module with basis $\{[D(\lambda)] \mid \lambda \in \mathcal{RP}_d\}$. For each $\lambda \in \mathcal{RP}_d$, let $P(\lambda)$ be the projective cover of $D(\lambda)$; in particular there exists a degree preserving surjection $P(\lambda) \twoheadrightarrow D(\lambda)$.

By definition of the graded decomposition numbers $d_{\mu, \lambda}$, for every $\mu \in \mathcal{P}_d$, we have $[S(\mu)] = \sum_{\lambda \in \mathcal{RP}_d} d_{\mu, \lambda} [D(\lambda)]$ in $[\text{Rep}(H_d)]$. Moreover, from [BK3, Theorem 3.14, Theorem 5.13] it follows that in $[\text{Rep}(H_d)]$ we have:

$$[P(\lambda)] = \sum_{\mu \in \mathcal{P}_d} d_{\mu, \lambda} [S(\mu)] \quad (\lambda \in \mathcal{RP}_d). \quad (5.1)$$

The *Fock space* $\mathfrak{F}$ is a $U_q(\mathfrak{sl}_e)$ -module with $\mathbb{Q}(q)$ -basis $\{\mu \mid \mu \in \bigoplus_{d \geq 0} \mathcal{P}_d\}$. The submodule of $\mathfrak{F}$ generated by the vector $\emptyset$ (corresponding to the empty partition) is the irreducible highest weight module $V(\Lambda_0)$. There is a canonical $U_q(\mathfrak{sl}_e)$ -module homomorphism $\pi : \mathfrak{F} \rightarrow V(\Lambda_0)$, see [BK3, (3.29)]. We can and always will identify $\bigoplus_{d \geq 0} [\text{Rep}(H_d)] \otimes_{\mathbb{Z}[q, q^{-1}]} \mathbb{Q}(q) = V(\Lambda_0) \subset \mathfrak{F}$. Under this identification, we have $\pi(\mu) = [S(\mu)]$ for each μ .

5.2. Projective H_d -modules. For each $\mathbf{i} \in I^d$ there is a unique idempotent $e(\mathbf{i}) \in H_d$ (possibly zero) such that $e(\mathbf{i})V = V_{\mathbf{i}}$ for any finite dimensional H_d -module, see e.g. [BK2].

Lemma 5.1. *If $\lambda \in \mathcal{RP}_d$, then $H_d e(\mathbf{j}^\lambda) = \bigoplus_{\mu \in \mathcal{RP}_d} m_\lambda(D(\mu)) P(\mu)$.*

Proof. If M and N are graded H_d -modules, we denote by $\text{HOM}_{H_d}(M, N)$ the graded vector space which consists of *all*, and not necessarily homogeneous, H_d -homomorphisms from M to N . Now, the graded multiplicity of $P(\mu)$ in $H_d e(\mathbf{j}^\lambda)$ is equal to

$$\text{qdim } \text{HOM}_{H_d}(H_d e(\mathbf{j}^\lambda), D(\mu)) = \text{qdim } e(\mathbf{j}^\lambda) D(\mu) = m_\lambda(D(\mu)),$$

as required. $\square$

Corollary 5.2. *For $\lambda \in \mathcal{RP}_d$, we have in $[\text{Rep}(H_d)]$:*

- (i) $[H_d e(\mathbf{j}^\lambda)] = \sum_{\mu \in \mathcal{RP}_d} m_\lambda(D(\mu)) [P(\mu)]$,
- (ii) $[H_d e(\mathbf{j}^\lambda)] = \sum_{\mu \in \mathcal{P}_d} m_\lambda(S(\mu)) [S(\mu)]$,
- (iii) $[H_d e(\mathbf{j}^\lambda)] = \sum_{\mu \in \mathcal{RP}_d} m_\lambda(P(\mu)) [D(\mu)]$.

Proof. (i) is a restatement of Lemma 5.1 in the Grothendieck group. (ii) follows from (i) and (5.1) since:

$$\begin{aligned} \sum_{\mu \in \mathcal{RP}_d} m_\lambda(D(\mu)) [P(\mu)] &= \sum_{\mu \in \mathcal{RP}_d} m_\lambda(D(\mu)) \sum_{\nu \in \mathcal{P}_d} d_{\nu, \mu} [S(\nu)] \\ &= \sum_{\nu \in \mathcal{P}_d} \left(\sum_{\mu \in \mathcal{RP}_d} m_\lambda(D(\mu)) d_{\nu, \mu} \right) [S(\nu)] = \sum_{\nu \in \mathcal{P}_d} m_\lambda(S(\nu)) [S(\nu)]. \end{aligned}$$

Similarly, (iii) follows from (ii) and (5.1) since:

$$\begin{aligned} \sum_{\mu \in \mathcal{P}_d} m_\lambda(S(\mu))[S(\mu)] &= \sum_{\mu \in \mathcal{P}_d} m_\lambda(S(\mu)) \sum_{\nu \in \mathcal{RP}_d} d_{\mu,\nu}[D(\nu)] \\ &= \sum_{\nu \in \mathcal{RP}_d} \left(\sum_{\mu \in \mathcal{P}_d} m_\lambda(S(\mu)) d_{\mu,\nu} \right) [D(\nu)] = \sum_{\nu \in \mathcal{RP}_d} m_\lambda(P(\nu))[S(\nu)]. \end{aligned}$$

This completes the proof. $\square$

The element $[H_d e(\mathbf{j}^\lambda)]$ is connected to the ‘first approximation’ $A(\lambda)$ from [LLT] as follows. It is easy to check that $A(\lambda) = \frac{1}{r_\lambda} \sum_{\mu \in \mathcal{P}_d} m_\lambda(S(\mu))\mu \in \mathfrak{F}$, and thus $\pi(r_\lambda A(\lambda)) = [H_d e(\mathbf{j}^\lambda)]$ using Corollary 5.2(ii). This gives an explicit connection between the formal sum $A(\lambda) \in \mathfrak{F}$ and the representation theory of H_d .

5.3. Comparing the algorithms. Consider the following system of equations over $\mathbb{Z}[q, q^{-1}]$:

$$m_\lambda(S(\mu)) = \sum_{\nu \in \mathcal{RP}_d} d_{\mu,\nu} m_\lambda(D(\nu)) \quad (\lambda \in \mathcal{RP}_d, \mu \in \mathcal{P}_d), \quad (5.2)$$

with unknowns $d_{\mu,\nu}$ and $m_\lambda(D(\nu))$ for $\mu \in \mathcal{P}_d$ and $\lambda, \nu \in \mathcal{RP}_d$. Note that $m_\lambda(S(\mu))$ are known from Theorem 2.2.

The algorithm described in section 4.2 allows us to solve the system (5.2) and relies on the fact that it has a unique solution under the following conditions:

- (i) all $m_\lambda(D(\nu)) \in \mathbb{Z}[q, q^{-1}]$ are bar-invariant and $m_\lambda(D(\lambda)) = r_\lambda$;
- (ii) $d_{\mu,\nu} = 0$ unless $\nu \leq \mu$, $d_{\nu,\nu} = 1$, and $d_{\mu,\nu} \in q\mathbb{Z}[q]$ for $\mu \neq \nu$.

It turns out that the LLT algorithm also relies on being able to solve the system (5.2). In [LLT], the vector $A(\lambda)$ for $\lambda \in \mathcal{RP}_d$ is a first approximation to the canonical basis element $G(\lambda)$, which calculates a *column* of the decomposition matrix: $G(\lambda) = \sum_{\mu \in \mathcal{P}_d} d_{\mu,\lambda} \mu$. One may write $r_\lambda A(\lambda) = \sum_{\nu \in \mathcal{RP}_d} b_{\lambda,\nu} G(\nu)$ for some bar-invariant coefficients $b_{\lambda,\nu}$. Then:

$$\sum_{\mu \in \mathcal{P}_d} m_\lambda(S(\mu))\mu = \sum_{\nu \in \mathcal{RP}_d} b_{\lambda,\nu} \sum_{\mu \in \mathcal{P}_d} d_{\mu,\nu} \mu.$$

Fixing μ (i.e. fixing a *row* of the matrix) we are left with solving

$$m_\lambda(S(\mu)) = \sum_{\nu \in \mathcal{RP}_d} b_{\lambda,\nu} d_{\mu,\nu}$$

for each $\mu \in \mathcal{P}_d$ and $\lambda \in \mathcal{RP}_d$. Since conditions analogous to (i) and (ii) are known to hold in the Lascoux-Leclerc-Thibon setup [LLT], we are left with solving the same system of equations under the same conditions. Moreover, $b_{\lambda,\nu}$ appearing in the LLT algorithm can now be interpreted as the graded weight space multiplicities $m_\lambda(D(\nu))$ for each $\lambda, \nu \in \mathcal{RP}_d$.

REFERENCES

- [A] S. Ariki, On the decomposition numbers of the Hecke algebra of $G(m, 1, n)$, *J. Math. Kyoto Univ.* **36** (1996), 789–808.
- [BK1] J. Brundan and A. S. Kleshchev, Representation Theory of Symmetric Groups and their Double Covers, in: *Groups, Combinatorics, and Geometry (Durham, 2001)*, pp.31–53, World Scientific Publishing, River Edge, NJ, 2003.
- [BK2] J. Brundan and A. S. Kleshchev, Blocks of cyclotomic Hecke algebras and Khovanov-Lauda algebras, *Invent. Math.*, to appear.
- [BK3] J. Brundan and A. S. Kleshchev, Graded decomposition numbers for cyclotomic Hecke algebras, *Adv. in Math.*, to appear.
- [BKW] J. Brundan, A. S. Kleshchev and W. Wang, Graded Specht Modules, [arXiv:0901.0218](#).
- [DJ] R. Dipper and G.D. James, Representations of Hecke algebras of general linear groups, *Proc. London Math Soc.*, **52** (1986), 20–52.
- [DJM] R. Dipper, G.D. James, and A. Mathas, Cyclotomic q -Schur algebras, *Math. Z.* **229** (1998), 385–416.
- [G] I. Grojnowski, Affine $\mathfrak{sl}_p$ controls the representation theory of the symmetric group and related Hecke algebras, [arXiv:math.RT/9907129](#)
- [J1] G.D. James, *The Representation Theory of the Symmetric Groups*, Lecture Notes in Mathematics 682, Springer, Berlin, 1978.
- [J2] G. D. James, On the decomposition matrices of the symmetric groups. II, *J. Algebra* **43** (1976), 45–54.
- [J3] G.D. James, Representations of the symmetric groups over the field of order 2, *J. Algebra* **38** (1976), 280–308.
- [K] A. Kleshchev, *Linear and Projective Representations of Symmetric Groups*, Cambridge University Press, Cambridge, 2005.
- [KL] M. Khovanov and A. Lauda, A diagrammatic approach to categorification of quantum groups I; [arXiv:0803.4121](#).
- [KS] A. S. Kleshchev and J. Sheth, On extensions of simple modules over symmetric and algebraic groups, *J. Algebra* **221** (1999), 705–722.
- [LLT] A. Lascoux, B. Leclerc, and J.-Y. Thibon, Hecke algebras at roots of unity and crystal bases of quantum affine algebras, *Commun. Math. Phys.* **181** (1996), 205–263.
- [L] S. Lyle, Some results obtained by application of the LLT algorithm, *Communications in Algebra* **34** (2006), 1723–1752.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, USA.

E-mail address: klesh@uoregon.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, USA.

E-mail address: dnash@uoregon.edu